

INSTABILITY OF ISOLATED SPECTRUM FOR W-SHAPED MAPS

ZHENYANG LI AND PAWEŁ GÓRA

ABSTRACT. In this note we consider W -shaped map $W_0 = W_{s_1, s_2}$ with $\frac{1}{s_1} + \frac{1}{s_2} = 1$ and show that eigenvalue 1 is not stable. We do this in a constructive way. For each perturbing map W_a we show the existence of the “second” eigenvalue λ_a , such that $\lambda_a \rightarrow 1$, as $a \rightarrow 0$, which proves instability of isolated spectrum of W_0 . At the same time, the existence of second eigenvalues close to 1 causes the maps W_a behave in a metastable way. They have two almost invariant sets and the system spends long periods of consecutive iterations in each of them with infrequent jumps from one to the other.

1. INTRODUCTION

One of the most important problems in the theory of dynamical systems is their stability and possible instability. In particular, in the theory of piecewise expanding maps of interval, it is interesting whether the given system has a stable absolutely continuous invariant measure (acim), and more generally, if the isolated spectrum of Perron-Frobenius operator is stable under small perturbations of the map. For a general introduction to the theory of piecewise expanding one-dimensional maps we refer the reader to [1]. Most relevant to the stability problems are papers [5] and [6].

In general, the setting of the stability problem we are interested in is as follows: Let τ_0 be a piecewise expanding map of an interval with unique acim μ_0 and $\{\tau_a\}_{a>0}$ a family of its perturbations with acims μ_a , correspondingly. If maps τ_a converge to τ_0 (say, in Skorokhod metric), do their acims converge (say, in $*$ -weak topology) to μ_0 ? Or more generally, do the isolated spectra of P_{τ_a} converge to the isolated spectrum of P_{τ_0} , including multiplicities and eigenfunctions? P_τ is the Perron-Frobenius operator induced by τ on the space of functions

Date: October 18, 2011.

2000 Mathematics Subject Classification. Primary 37A10, 37A05, 37E05.

Key words and phrases. piecewise expanding maps, Frobenius-Perron operator, Markov maps, W -shaped maps, second eigenvalue, stability, metastable behaviour.

The research of the authors was supported by NSERC grants.

of bounded variation and by isolated spectrum we mean the part of the spectrum which lies outside the essential spectral radius. Papers [5] and [6] show that such stability takes place if the family $\{\tau_a\}_{a \geq 0}$ satisfies Lasota-Yorke inequality ([7] or [2] for strengthened form) with uniform constants. Usual conditions ensuring this are $|\tau'_a| > 2 + \varepsilon$ plus the minimal length of subintervals of defining partitions uniformly separated from 0.

One of the known sources of instability is the presence of turning fixed or periodic point touching a map branch with slope 2 or smaller. The famous example is the W -shaped map introduced in [5]. Because of the turning fixed point we cannot use an iterate of the map to increase the minimal slope. It causes appearance of arbitrary short partition intervals in perturbed maps.

Recently the interest in W -shaped maps increased. We will introduce them in more detail. The W_{s_1, s_2} map is a piecewise linear map of the interval $[0, 1]$ onto itself with a graph in the shape of letter W . The first and the third branch are decreasing, the second and the last increasing. The first and the last branches are onto and with relatively large slopes (usually around -4 and 4). The second branch has slope s_1 and the third has slope $-s_2$ and $1/2$ is the point where they meet. $W_{s_1, s_2}(1/2) = 1/2$ so $1/2$ is the turning fixed point. The original W -map of [5] is of $W_{2,2}$ type and it was proved there that its acim is unstable under some family of localized perturbations. In [3] a family of non-local Markov perturbations was constructed which also caused the instability of acim of $W_{2,2}$. By non-local we mean that each perturbed map is exact on the whole $[0, 1]$. This result has been generalized in [8] where a whole continuous family of perturbations was constructed with the same effect. More general situation was considered in [9]. The perturbations similar to that of [8] were considered. It was shown there that depending on whether $\frac{1}{s_1} + \frac{1}{s_2}$ is larger, equal or smaller than 1 the limit of μ_a 's is Dirac measure $\delta_{1/2}$, or a combination of $\delta_{1/2}$ and μ_0 , or μ_0 , correspondingly. This result suggested that condition $\frac{1}{s_1} + \frac{1}{s_2} < 1$ may actually imply stability which was later proved for a quite general setting in [2].

In this note we consider map W_0 of type W_{s_1, s_2} with $\frac{1}{s_1} + \frac{1}{s_2} = 1$ and show that that eigenvalue 1 is not stable. We do this in a constructive way. For each perturbed map W_a we show the existence of the “second” eigenvalue λ_a ,

$$\frac{1 - 2ra}{1 + 2ra} < \lambda_a < \frac{1}{1 + 2ra} ,$$

where r is a constant independent of a . Thus, as $a \rightarrow 0$ the eigenvalues $\lambda_a \rightarrow 1$ which shows instability of isolated spectrum of W_0 . At the same time, the existence of second eigenvalues close to 1 causes the maps W_a behave in a metastable way. They have two almost invariant sets and the system spends long periods of consecutive iterations in each of them with infrequent jumps from one to the other.

2. MARKOV W_a MAPS AND THEIR INVARIANT DENSITIES

Let $s_1, s_2 > 1$ satisfy $\frac{1}{s_1} + \frac{1}{s_2} = 1$, and $r > 0$. Let us consider W -shaped map:

$$W_0(x) = \begin{cases} W_{0,1}(x) := 1 - 2s_2x, & 0 \leq x < \frac{1}{2} - \frac{1}{2s_1}, \\ W_{0,2}(x) := s_1(x - \frac{1}{2} + \frac{1}{2s_1}), & \frac{1}{2} - \frac{1}{2s_1} \leq x < \frac{1}{2}, \\ W_{0,3}(x) := s_2(\frac{1}{2} + \frac{1}{2s_2} - x), & \frac{1}{2} \leq x < \frac{1}{2} + \frac{1}{2s_2}, \\ W_{0,4}(x) := 2s_1(x - 1) + 1, & \frac{1}{2} + \frac{1}{2s_2} \leq x < 1, \end{cases}$$

and its perturbations, W_a maps with parameter $a > 0$:

$$W_a(x) = \begin{cases} W_{a,1}(x) := 1 - 2s_2x, & 0 \leq x < \frac{1}{2} - \frac{1}{2s_1}, \\ W_{a,2}(x) := (s_1 + 2rs_1a)(x - \frac{1}{2} + \frac{1}{2s_1}), & \frac{1}{2} - \frac{1}{2s_1} \leq x < \frac{1}{2}, \\ W_{a,3}(x) := (s_2 + 2rs_2a)(\frac{1}{2} + \frac{1}{2s_2} - x), & \frac{1}{2} \leq x < \frac{1}{2} + \frac{1}{2s_2}, \\ W_{a,4}(x) := 2s_1(x - 1) + 1, & \frac{1}{2} + \frac{1}{2s_2} \leq x < 1. \end{cases}$$

Let $\tau_i = W_{a,i}^{-1}$, $i = 1, 2, 3, 4$; $I_0 = [0, \frac{1}{2} + ra]$. The Frobenius-Perron operator (see [1]) associated with W_a is

$$\begin{aligned} P_a f = \frac{1}{2s_2} f \circ \tau_1 + \frac{1}{s_1 + 2rs_1a} (f \circ \tau_2) \chi_{I_0} &+ \frac{1}{s_2 + 2rs_2a} (f \circ \tau_3) \chi_{I_0} \\ &+ \frac{1}{2s_1} f \circ \tau_4 \end{aligned}$$

Note that

$$(1) \quad \begin{aligned} \chi_{I_0} \circ \tau_1 &= 1, \quad \chi_{I_0} \circ \tau_2 = \chi_{I_0}, \\ \chi_{I_0} \circ \tau_3 &= \chi_{[W_a^2(1/2), \frac{1}{2} + ra]}, \quad \chi_{I_0} \circ \tau_4 = 0. \end{aligned}$$

Let $I_1 = [W_a^2(1/2), \frac{1}{2} + ra]$ whose left end point is $W_a^2(\frac{1}{2}) = W_a(\frac{1}{2} + ra)$.

We will consider only parameters a such that W_a is a Markov map, i.e., some iterate of $1/2$ falls into an endpoint of the defining partition. Let a satisfy:

$$(2) \quad W_a^m(\frac{1}{2} + ra) = \frac{1}{2} - \frac{1}{2s_1},$$

where $m \geq 1$ is the first time when the trajectory of $W_a(\frac{1}{2}) = \frac{1}{2} + ra$ reaches the partition point $\frac{1}{2} - \frac{1}{2s_1}$. Note that $\frac{1}{2} - \frac{1}{2s_1} = \frac{1}{2s_2}$. The point

$W_a(\frac{1}{2} + ra)$ is just below the fixed point on the second branch of W_a and the consecutive images $W_a^i(\frac{1}{2} + ra)$ decrease until for $i = m$ the equality (2) is satisfied.

Let us take 1 as the initial function and iterate it using operator P_a . Let $P_a^n 1$ be denoted by $f_{n,m}$. Let

$$I_i = [W_a^i(\frac{1}{2} + ra), \frac{1}{2} + ra], i = 1, 2, \dots, m.$$

Because of (2) and (1), after some number of iterations ($n \geq m + 1$), we have:

$$f_{n,m} = c_{n,0} + \alpha_{n,0}\chi_{I_0} + \alpha_{n,1}\chi_{I_1} + \alpha_{n,2}\chi_{I_2} + \dots + \alpha_{n,m-1}\chi_{I_{m-1}} + \alpha_{n,m}\chi_{I_m},$$

where $c_{n,0}$ and $\alpha_{n,i}$ ($i = 0, 1, \dots, m$) are constants. Now, let us look at the $f_{n+1,m}$. We have the following proposition.

Proposition 1. (I) $c_{n,0} \circ \tau_1$ and $c_{n,0} \circ \tau_4$ are again constant functions, $c_{n,0} \circ \tau_2\chi_{I_0}$ and $c_{n,0} \circ \tau_3\chi_{I_0}$ are the characteristic function χ_{I_0} ;

(II) $\chi_{I_0} \circ \tau_1$ is constant function, $\chi_{I_0} \circ \tau_2\chi_{I_0} = \chi_{I_0}$, $\chi_{I_0} \circ \tau_3\chi_{I_0} = \chi_{I_1}$, $\chi_{I_0} \circ \tau_4$ is 0;

(III) For $i = 1, 2, \dots, m-1$, $\chi_{I_i} \circ \tau_1$ and $\chi_{I_i} \circ \tau_4$ are 0, $\chi_{I_i} \circ \tau_2\chi_{I_0} = \chi_{I_{i+1}}$, $\chi_{I_i} \circ \tau_3\chi_{I_0} = \chi_{I_1}$;

(IV) $\chi_{I_m} \circ \tau_1$ and $\chi_{I_m} \circ \tau_4$ are 0, $\chi_{I_m} \circ \tau_2\chi_{I_0} = \chi_{I_0}$, $\chi_{I_m} \circ \tau_3\chi_{I_0} = \chi_{I_1}$.

Thus, we have the following proposition.

Proposition 2. for n big enough, $f_{n,m}$ always has the form:

$$f_{n,m} = c_{n,0} + \alpha_{n,0}\chi_{I_0} + \alpha_{n,1}\chi_{I_1} + \alpha_{n,2}\chi_{I_2} + \dots + \alpha_{n,m-1}\chi_{I_{m-1}} + \alpha_{n,m}\chi_{I_m},$$

and

$$\begin{bmatrix} c_{n+1,0} \\ \alpha_{n+1,0} \\ \alpha_{n+1,1} \\ \vdots \\ \alpha_{n+1,m} \end{bmatrix} = A_m \begin{bmatrix} c_{n,0} \\ \alpha_{n,0} \\ \alpha_{n,1} \\ \vdots \\ \alpha_{n,m} \end{bmatrix}.$$

where $(m+2) \times (m+2)$ matrix A_m is given by

$$A_m = \begin{bmatrix} \frac{1}{2s_1} + \frac{1}{2s_1} & \frac{1}{2s_2} & 0 & 0 & 0 & \cdots & 0 & 0 \\ \frac{1}{s_1+2rs_1a} + \frac{1}{s_2+2rs_2a} & \frac{1}{s_1+2rs_1a} & 0 & 0 & 0 & \cdots & 0 & \frac{1}{s_1+2rs_1a} \\ 0 & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} & \cdots & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} \\ 0 & 0 & \frac{1}{s_1+2rs_1a} & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{s_1+2rs_1a} & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \cdots & \frac{1}{s_1+2rs_1a} & 0 \end{bmatrix}.$$

Since $\frac{1}{s_1} + \frac{1}{s_2} = 1$, we can simplify the A_m to the form

$$A_m = \begin{bmatrix} \frac{1}{2} & \frac{1}{2s_2} & 0 & 0 & 0 & \cdots & 0 & 0 \\ \frac{1}{1+2ra} & \frac{1}{s_1+2rs_1a} & 0 & 0 & 0 & \cdots & 0 & \frac{1}{s_1+2rs_1a} \\ 0 & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} & \cdots & \frac{1}{s_2+2rs_2a} & \frac{1}{s_2+2rs_2a} \\ 0 & 0 & \frac{1}{s_1+2rs_1a} & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{s_1+2rs_1a} & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \cdots & \frac{1}{s_1+2rs_1a} & 0 \end{bmatrix}.$$

We also need the following proposition.

Proposition 3. Equation (2) is equivalent to:

$$(s_2 + 2rs_2a)(s_1 + 2rs_1a)^{m-1} - \sum_{i=0}^{m-1} (s_1 + 2rs_1a)^i = \frac{1}{2rs_1a},$$

or

$$(3) \quad (s_1 + 2rs_1a)^m = \frac{1}{4r^2s_2^2a^2}.$$

Proof. If

$$(s_2 + 2rs_2a)(s_1 + 2rs_1a)^{m-1} - \sum_{i=0}^{m-1} (s_1 + 2rs_1a)^i = \frac{1}{2rs_1a},$$

then

$$(s_1+2rs_1a)^{m-1} [(s_2 + 2rs_2a)(s_1 + 2rs_1a - 1) - (s_1 + 2rs_1a)] = \frac{s_1 - 1}{2rs_1a} = \frac{1}{2rs_2a},$$

Thus, we obtain

$$(s_1 + 2rs_1a)^m = \frac{1}{4r^2s_2^2a^2}.$$

On the other hand, it is proven in [9] that

$$\begin{aligned} W_a^{m+1}(1/2) \\ = -a^2(s_1 + 2rs_1a)^{m-1} \frac{r(2rs_1s_2 + 2rs_1s_2 - 2rs_1 - 2rs_2) + 4r^3s_1s_2a}{s_1 + 2rs_1a - 1} \\ + \frac{s_1 - 1 + 2rs_1a - 2ra}{2(s_1 + 2rs_1a - 1)}. \end{aligned}$$

If equation (2) holds, then

$$a^2(s_1 + 2rs_1a)^{m-1} \frac{2r^2s_1s_2 + 4r^3s_1s_2a}{s_1 + 2rs_1a - 1} = \frac{s_1 - 1}{2s_1(s_1 + 2rs_1a - 1)},$$

hence,

$$(s_1 + 2rs_1a)^{m-1} = \frac{1}{a^24r^2s_1s_2^2(1 + 2ra)}$$

which is equivalent to equation (3). $\square$

Using Proposition 2, we can find the fixed vector of A_m . Let us denote it by $(c, \alpha_0, \alpha_1, \dots, \alpha_m)^T$. Then, the fixed function (not necessarily normalized) of P_a is:

$$g_m^* = c + \alpha_0\chi_{I_0} + \alpha_1\chi_{I_1} + \alpha_2\chi_{I_2} + \dots + \alpha_{m-1}\chi_{I_{m-1}} + \alpha_m\chi_{I_m},$$

where

$$\begin{aligned} c &= \frac{1}{2rs_1s_2a} \\ \alpha_0 &= \frac{1}{2rs_1a} \\ \alpha_1 &= (s_1 + 2rs_1a)^{m-1} \\ \alpha_2 &= (s_1 + 2rs_1a)^{m-2} \\ &\dots \\ \alpha_{m-2} &= (s_1 + 2rs_1a)^2 \\ \alpha_{m-1} &= s_1 + 2rs_1a \\ \alpha_m &= 1. \end{aligned}$$

It was proven in [9] that after normalization the measures $g_m^* \cdot L$ converge to the measure

$$\frac{1}{2r(s_1 + s_2)(s_2 + 2) + 2rs_1s_2^2} (2r(s_1 + s_2)(s_2 + 2)\mu_0 + 2rs_1s_2^2\delta_{1/2}) ,$$

where L is Lebesgue measure, μ_0 is the absolutely continuous invariant measure of W_0 and $\delta_{1/2}$ is Dirac measure at $1/2$.

3. SECOND EIGENVALUES FOR MARKOV W_a MAPS

Now, instead for a fixed vector, we will look for an eigenvector corresponding to an eigenvalue $\lambda < 1$. Denote the eigenvector of A_m associated with λ by $(c, \alpha_0, \alpha_1, \dots, \alpha_m)^T$. Then, the corresponding eigenfunction of P_a associated with λ is:

$$(4) \quad h_m = c + \alpha_0 \chi_{I_0} + \alpha_1 \chi_{I_1} + \alpha_2 \chi_{I_2} + \dots + \alpha_{m-1} \chi_{I_{m-1}} + \alpha_m \chi_{I_m} .$$

The equation

$$A_m h_m = \lambda h_m ,$$

is equivalent to the system

$$\begin{aligned} \lambda c &= \frac{1}{2}c + \frac{1}{2s_2}\alpha_0 \\ \lambda \alpha_0 &= \frac{1}{1+2ra}\left(c + \frac{1}{s_1}\alpha_0 + \frac{1}{s_1}\alpha_m\right) \\ \lambda \alpha_1 &= \frac{1}{s_2(1+2ra)}(\alpha_0 + \alpha_1 + \dots + \alpha_m) \\ \lambda \alpha_2 &= \frac{1}{s_1(1+2ra)}\alpha_1 \\ \lambda \alpha_3 &= \frac{1}{s_1(1+2ra)}\alpha_2 \\ &\dots \\ \lambda \alpha_{m-2} &= \frac{1}{s_1(1+2ra)}\alpha_{m-3} \\ \lambda \alpha_{m-1} &= \frac{1}{s_1(1+2ra)}\alpha_{m-2} \\ \lambda \alpha_m &= \frac{1}{s_1(1+2ra)}\alpha_{m-1}. \end{aligned}$$

We can solve this system starting from the last equation. Let $\alpha_m = 1$. Then,

$$\begin{aligned}
 (5) \quad & \alpha_m = 1 \\
 & \alpha_{m-1} = \lambda s_1(1 + 2ra) \\
 & \alpha_{m-2} = \lambda^2 s_1^2(1 + 2ra)^2 \\
 & \dots \\
 & \alpha_2 = \lambda^{m-2} s_1^{m-2}(1 + 2ra)^{m-2} \\
 & \alpha_1 = \lambda^{m-1} s_1^{m-1}(1 + 2ra)^{m-1} \\
 & \alpha_0 = \lambda s_2(1 + 2ra)\alpha_1 - (\alpha_1 + \alpha_2 + \dots + \alpha_m) \\
 & = \lambda^m s_1^{m-1} s_2(1 + 2ra)^m - \frac{\lambda^m s_1^m(1 + 2ra)^m - 1}{\lambda s_1(1 + 2ra) - 1} \\
 & = \frac{\lambda^m s_1^{m-1}(1 + 2ra)^m(\lambda s_1 s_2(1 + 2ra) - s_1 s_2) + 1}{\lambda s_1(1 + 2ra) - 1} \\
 & c = \lambda(1 + 2ra)\alpha_0 - \frac{\alpha_0}{s_1} - \frac{1}{s_1} \\
 & c = \frac{\alpha_0}{s_2(2\lambda - 1)}
 \end{aligned}$$

We have two expressions for c . The system can be solved only if they are equal. Thus, we obtain equation

$$\begin{aligned}
 & \lambda^m s_1^{m-2}(1 + 2ra)^m(\lambda s_1 s_2(1 + 2ra) - s_1 s_2) \\
 & = \frac{\lambda^m s_1^{m-1}(1 + 2ra)^m(\lambda s_1 s_2(1 + 2ra) - s_1 s_2) + 1}{s_2(2\lambda - 1)(\lambda s_1(1 + 2ra) - 1)},
 \end{aligned}$$

or

$$\lambda^m s_1^{m-1} s_2(1 + 2ra)^m(\lambda(1 + 2ra) - 1)[s_2(2\lambda - 1)(\lambda s_1(1 + 2ra) - 1) - s_1] = 1.$$

We are going to prove that for small a this equation has a solution $\frac{1-2ra}{1+2ra} < \lambda < \frac{1}{1+2ra}$. Let us introduce an auxiliary function

$$\phi(\lambda) = \lambda^m s_1^{m-1} s_2(1 + 2ra)^m(\lambda(1 + 2ra) - 1)[s_2(2\lambda - 1)(\lambda s_1(1 + 2ra) - 1) - s_1].$$

Obviously $\phi(\frac{1}{1+2ra}) = 0$. We will show that $\phi(\frac{1-2ra}{1+2ra}) > 1$ if a is small enough. We have

$$\begin{aligned}
 \phi\left(\frac{1-2ra}{1+2ra}\right) &= \left(\frac{1-2ra}{1+2ra}\right)^m s_1^{m-1} s_2(1 + 2ra)^m \left(\left(\frac{1-2ra}{1+2ra}\right)(1 + 2ra) - 1 \right) \\
 &\quad \cdot \left[s_2 \left(2 \left(\frac{1-2ra}{1+2ra}\right) - 1 \right) \left(\left(\frac{1-2ra}{1+2ra}\right) s_1(1 + 2ra) - 1 \right) - s_1 \right] \\
 &= (1 - 2ra)^m s_1^{m-1} s_2(-2ra) \frac{-2ra(s_2 + 5s_1 - 6s_1 s_2 ra)}{1 + 2ra}
 \end{aligned}$$

$$= (1 - 2ra)^m s_1^{m-1} s_2 4r^2 a^2 \frac{s_2 + 5s_1 - 6s_1 s_2 ra}{1 + 2ra}.$$

Using (3) we obtain

$$\begin{aligned} \phi\left(\frac{1 - 2ra}{1 + 2ra}\right) &= \left(\frac{1 - 2ra}{1 + 2ra}\right)^m \frac{s_2 + 5s_1 - 6s_1 s_2 ra}{s_1 s_2 (1 + 2ra)} \\ &= \left(\frac{1 - 2ra}{1 + 2ra}\right)^m \frac{1 + \frac{4}{s_2} - 6ra}{1 + 2ra}. \end{aligned}$$

Note that if $a < \frac{1}{2rs_2}$, then $\frac{1 + \frac{4}{s_2} - 6ra}{1 + 2ra} > 1$. Furthermore, $\lim_{a \rightarrow 0} \frac{1 + \frac{4}{s_2} - 6ra}{1 + 2ra} = 1 + \frac{4}{s_2} > 1$.

Using (3) again, we can represent m as

$$(6) \quad m = \frac{-2 \ln(2s_2 ra)}{\ln(s_1 + 2s_1 ra)},$$

which gives

$$\begin{aligned} \phi\left(\frac{1 - 2ra}{1 + 2ra}\right) &= \frac{1 + \frac{4}{s_2} - 6ra}{1 + 2ra} \left(\frac{1 - 2ra}{1 + 2ra}\right)^{\frac{-2 \ln(2s_2 ra)}{\ln(s_1 + 2s_1 ra)}} \\ &= \frac{1 + \frac{4}{s_2} - 6ra}{1 + 2ra} \exp\left(-2 \ln\left(\frac{1 - 2ra}{1 + 2ra}\right) \frac{\ln(2s_2 ra)}{\ln(s_1 + 2s_1 ra)}\right). \end{aligned}$$

Since $\lim_{a \rightarrow 0} \ln\left(\frac{1 - 2ra}{1 + 2ra}\right) \ln(a) = 0$, the argument of exp converges to 0 as $a \rightarrow 0$. Thus

$$\lim_{a \rightarrow 0} \phi\left(\frac{1 - 2ra}{1 + 2ra}\right) = 1 + \frac{4}{s_2}.$$

This proves our claim for a small enough. We proved the following

Theorem 1. *Assume that a satisfies (3), for some integer m , i.e., W_a map is Markov with $W_a^{m+1}(1/2) = \frac{1}{2} - \frac{1}{2s_1}$. For a small enough, Perron-Frobenius operator P_a has an eigenvalue λ_a satisfying*

$$(7) \quad \frac{1 - 2ra}{1 + 2ra} < \lambda_a < \frac{1}{1 + 2ra}.$$

The corresponding eigenfunction is given by equations (4) and (5), up to a multiplicative constant.

Remark 1. *Using tedious calculations we were able to show that ϕ'' is positive in a neighbourhood of 1. Since $\phi\left(\frac{1 - 2ra}{1 + 2ra}\right) > 1$, $\phi\left(\frac{1}{1 + 2ra}\right) = 0$ and $\phi(1) = 1$, for small a there is only one eigenvalue in the interval $\left(\frac{1 - 2ra}{1 + 2ra}, 1\right)$, i.e., λ_a we found in Theorem 1 is really the “second” eigenvalue.*

4. EIGENFUNCTION FOR $\lambda_a < 1$

In this section we take a closer look at the eigenfunction corresponding to the second eigenvalue λ_a found in Theorem 1. We omit the subscript “a” to simplify the notation. Let $(c, \alpha_0, \alpha_1, \dots, \alpha_m)$ be the λ -eigenvector given by (5). We have

$$\alpha_j = \lambda^{m-j} s_1^{m-j} (1 + 2ra)^{m-j} > 0 \quad , \quad j = 1, \dots, m .$$

Next,

$$\alpha_0 = \frac{\lambda^m s_1^{m-1} (1 + 2ra)^m (\lambda s_1 s_2 (1 + 2ra) - s_1 s_2) + 1}{\lambda s_1 (1 + 2ra) - 1} < 0 ,$$

since $\lambda(1 + 2ra) < 1$ but very close to 1 and using formula (6) we can show that $\lambda^m (1 + 2ra)^m$ approaches 1 as $m \rightarrow \infty$. As $\alpha_0 < 0$, we also have

$$c = \frac{\alpha_0}{s_2(2\lambda - 1)} < 0 .$$

The P_a eigenfunction h_m , defined in (4), is positive on some interval $G_m = [W_a^{m_1}(1/2), 1/2 + a/4]$ and negative outside this interval. Since, as a decreases, more and more of numbers $\alpha_m, \alpha_{m-1}, \alpha_{m-2} \dots$ are necessary to balance $\alpha_0 + c$, we have $\lim_{a \rightarrow 0} (m - m_1) = +\infty$. This implies, that intervals G_m shrink to the point $1/2$ as $a \rightarrow 0$.

Since $0 < \lambda < 1$ we have $\int_0^1 h_m dL = 0$. Let $K_m = \int_0^1 |h_m| dL$. The normalized signed measures $\frac{1}{K_m} h_m \cdot L$ converge $*$ -weakly to the measure

$$-\frac{1}{2}\mu_0 + \frac{1}{2}\delta_{(1/2)} ,$$


where μ_0 is W_0 -invariant absolutely continuous measure and $\delta_{(1/2)}$ is Dirac measure at point $1/2$.

As it is described in [4] the presence of the eigenvalue λ close to 1 makes the system behave in a metastable way. The sets $A^+ = \{t : h_m(t) \geq 0\}$ and $A^- = \{t : h_m(t) < 0\}$ are almost invariant with the escape rates bounded by $-\ln \lambda$ which is close to 0. This means that a typical trajectory stays for a long time in A^+ , then jumps to A^- , stays there for a long time, then jumps to A^+ , spend there long time, etc. Despite the small essential spectral radius (equal to $\max\{1/s_1, 1/s_2\}$), the system converges to equilibrium slowly at the rate given by $C\lambda^n$, for some constant C .

Fig. 1 shows graphs of normalized functions h_m produced using Maple 13. We used $s_1 = s_2 = 2$ and $r = 1/4$.

a) $m = 5$, $a = 0.14789903570478$, $\lambda = 0.8732372308$, $K_m = 3.819456626$;

b) $m = 7$, $a = 0.077390319202550$, $\lambda = 0.9365803433$, $K_m = 8.987509817$.

FIGURE 1. Normalized eigenfunctions h_m .

Note that the vertical scales of the pictures are very different.

REFERENCES

- [1] A. Boyarsky and P. Góra, *Laws of Chaos. Invariant Measures and Dynamical Systems in One Dimension*, Probability and its Applications, Birkhäuser, Boston, MA, 1997.
- [2] P. Eslami and P. Góra, *Stronger Lasota-Yorke inequality for piecewise monotonic transformations*, preprint.
- [3] P. Eslami and M. Misiurewicz, *Singular limits of absolutely continuous invariant measures for families of transitive map*, Journal of Difference Equations and Applications, doi:10.1080/10236198.2011.590480.
- [4] G. Froyland and O. Stančević, *Escape rates and Perron-Frobenius operators: Open and closed dynamical systems*, Discrete and Continuous Dynamical Systems - Series B **14** (2010), 457–472.
- [5] G. Keller, *Stochastic stability in some chaotic dynamical systems*, Monatshefte für Mathematik **94** (4) (1982), 313–333.
- [6] G. Keller and C. Liverani., *Stability of the spectrum for transfer operators*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), **28** (1)(1999), 141–152.
- [7] A. Lasota; J. A. Yorke, *On the existence of invariant measures for piecewise monotonic transformations*, Trans. Amer. Math. Soc. **186** (1973), 481–488 (1974); MR0335758 (49 #538).
- [8] Z. Li, P. Góra, A. Boyarsky, H. Proppe and P. Eslami, *A family of piecewise expanding maps having singular measure as a limit of acim's*, in press, Ergodic Theory and Dynamical Systems.
- [9] Z. Li, *W-like maps with various instabilities of acim's*, available at <http://arxiv.org/abs/1109.5199>

DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY,
1455 DE MAISONNEUVE BLVD. WEST, MONTREAL, QUEBEC H3G 1M8, CANADA

E-mail address: `zhenyangemail@gmail.com`

E-mail address: `pgora@mathstat.concordia.ca`